

ABOUT US

Safecare Industries located in KEZAD- Abu Dhabi with fully automated manufacturing facilities specialized in manufacturing and supplying an extensive range of Microbiological Pre-poured Culture Media (PPM), Pre-poured Tubed Media and Bottle Media. All operational functions adhere to the required industry compliance criteria.

MISSION

Our mission is to Manufacturer and supply high quality Readymade Culture Media and broth and supply across UAE and GCC countries.

VISION

Our vision to provide high quality cost effective complete range of Microbiological Prepared Culture Media to UAE and GCC Market.

IMPORTANT SECTORS

Clinical, Veterinary Pharmaceutical, Industrial and Cosmetic

KEY FEATURES

- All our PPM products are quality assures and manufacturing facilities adhere to Good Manufacturing Practice Guidelines.
- We produce our PPM products using high quality dehydrated culture media/ supplements and automated production lines to get consistent performance.
- Safecare follows the stringent stands of Good Manufacturing Practices (GMP) defined and approved by Ministry of Health UAE. Class 10,000 validated clean area Fully automated 6 line production lines

- Fully automatic integrated online quality inspection camera vision system for capturing the images of the dishes and high speed expulsion station for extracting the not suitable dishes.
- Warehouse logistics function and warehouse located in Abu Dhabi, Dubai, Sharjah Emirates with Refrigerated vehicle facilitating the timeframe delivery.

ACCREDITION

- ISO 9001: 2015 Quality Management System
- ISO 13485: 2016 The development, manufacture and supply of ready-to-use culture media
- CE (European Conformity)
- International Good Manufacturing Practices (GMP) which is certified by Ministry of Health UAE.
- **MOH License**

QUALITY

Safecare Prepared plates and broth tested in accordance with CLSI M22 A3- Quality assurance for Commercially Prepared Microbiological Culture Media - second edition.

PRODUCTS

Prepared plates media 90 MM Prepared plates media 150 MM Prepared bioplates media 90 MM Prepared tube media Prepared broth media Prepared media in Bottles

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C) |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|------------|----------------------|----------------------|-----------------|
| Anaerobic Isolation Agar | MBN0105 | 10 x 90 | 60 | 2-8°C |
| A medium for the growth of anaerobic micro-organisms, | | mm plates | | |
| particularly Bacteroides spp. and other he aerobes. | | | | |
| Azide Sheep Blood agar | MBN0109 | 10 x 90 | 60 | 2-8℃ |
| A selective medium for the detection and isolation of | | mm plates | | |
| streptococci and staphylococci from faeces, sewage and other | | | | |
| specimens | | | | |
| Bacillus Cereus Selective Agar(MYP) | MBN0120 | 10 x 90 | 60 | 2-8℃ |
| A selective and diagnostic medium for the isolation and | | mm plates | | |
| enumeration of Bacillus cereus | | ' | | |
| Baird Parker Agar | MBN0123 | 10 x 90 | 60 | 2-8℃ |
| A selective and diagnostic medium for the isolation and | | mm plates | | |
| enumeration of coagulase-positive staphylococci in foods. | | mm places | | |
| BCYE GVPC agar | MBN0135 | 10 x 90 | 60 | 2-8℃ |
| Buffered Charcoal Yeast Extract Agar for the selective isolation of Legionella when used with Legionella BCYE Growth Supplement and Selective supplement | WENTO | mm plates | 00 | 200 |
| BCYE cysteine agar | MBN0136 | 10 x 90 | 60 | 2-8℃ |
| Charcoal Yeast Extract Agar for the isolation of Legionella when | IVIDINUTSO | | 00 | 2-0 C |
| used with Legionella BCYE Growth Supplement with L-cysteine | | mm plates | | |
| BCYE Agar without cysteine | MBN0137 | 10 x 90 | 60 | 2-8℃ |
| Charcoal Yeast Extract Agar for the isolation of Legionella when used with Legionella BCYE Growth Supplement without L-cysteine | | mm plates | | |
| Brain Heart Infusion (BHI) Agar | MBN0141 | 10 x 90 | 60 | 2-8°C |
| A versatile medium for the culture of fastidious organisms | | mm plates | | |
| Bordet Gengou Blood Agar | MBN0145 | 10 x 90 | 60 | 2-8℃ |
| For the detection and isolation of Bordetella pertussis and Bordetella parapertussis from clinical specimens | | mm plates | | |
| Brilliant Green Agar Highly selective medium for the isolation of Salmonella species | MBN0147 | 10 x 90 mm plates | 90 | 2-8℃ |
| other than Salmonella typhi from foods, faeces and dairy | | • | | |
| products | | | | |
| Brilliant Green Bile Agar | MBN0148 | 10 x 90 | 90 | 2-8℃ |
| For the determination of the degree of contamination by | | mm plates | | |
| coliforms in drinking water and wastewater | 1,000 | | | |
| Bile Esculin Azide Agar | MBN0149 | 10 x 90 | 60 | 2-8℃ |
| A Selective medium for the isolation and presumptive | | mm plates | | |
| identification of intestinal enterococci | MDNIO450 | 10 × 00 | 60 | 2.000 |
| Bile Esculin Agar For the isolation and presumptive identification of Enterococci /Group D Streptococci. | MBN0150 | 10 x 90 mm plates | 60 | 2-8℃ |
| Brucella Blood Agar | MBN0152 | 10 x 90 | 60 | 2-8℃ |
| Brucella blood agar with 5%sheep blood is used for the | | mm plates | | |
| qualitative isolation of nonfastidious microorganisms from a | | | | |
| | | | | |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(|
|-----------------------------------------------------------------|----------|-----------|----------------------|---------------|
| Campy 5 Blood Agar | MBN0170 | 10 x 90 | 60 | 2-8℃ |
| Compylobacter agar with 5 antimicrobics, 10%sheep blood is | | mm plates | | |
| recommended as a selective medium for the primary isolation | | | | |
| of Campylobacter jejuni from clinical specimens. | | | | |
| Campy Blood Free Agar | MBN0171 | 10 x 90 | 60 | 2-8℃ |
| A Selective medium for the isolation of Campylobacter jejuni, | | mm plates | | |
| Campylobacter coli and Campylobacter laridis | | | | |
| Campy Laked Blood Agar | MBN0172 | 10 x 90 | 60 | 2-8℃ |
| A Selective medium for the isolation of Campylobacter jejuni, | | mm plates | | |
| Campylobacter coli from human, animal, avian and | | | | |
| environmental specimens | | | | |
| Cetrimide agar | MBN0180 | 10 x 90 | 60 | 2-8℃ |
| Cetrimide agar is a selective medium for the isolation and | | mm plates | | |
| detection of <i>Pseudomonas aeruginosa</i> from pharmaceutical, | | | | |
| clinical and cosmetic samples. | | | | |
| Chocolate Agar | MBN0185 | 10 x 90 | 60 | 2-8°C |
| A medium for the isolation and cultivation of fastidious | | mm plates | | |
| microorganisms especially Neisseria and Haemophilus species | | | | |
| Chocolate Bacitracin Agar | MBN0186 | 10 x 90 | 60 | 2-8℃ |
| A highly nutritious medium enriched with horse blood, suitable | | mm plates | | |
| for the selective isolation of Haemophilus species | | | | |
| Clostridium Difficile Agar | MBN0189 | 10 x 90 | 60 | 2-8℃ |
| Clostridium difficile agar, when supplemented, is used for the | | mm plates | | |
| isolation of C. difficile from sample. | | | | |
| CLED Agar | MBN0192 | 10 x 90 | 90 | 2-8℃ |
| A non-selective differential plating medium for the growth and | | mm plates | | |
| enumeration of urinary tract microorganisms. | | | | |
| Columbia CNA Blood Agar | MBN0195 | 10 x 90 | 60 | 2-8℃ |
| A medium recommended for the isolation of gram positive | | mm plates | | |
| cocci especially staphylococci and streptococci from clinical | | | | |
| samples | | | | |
| Corn Meal Agar | MBN0196 | 10 x 90 | 90 | 2-8℃ |
| A recommended medium for chlamydospore production by | | mm plates | | |
| Candida albicans and for the maintenance of fungal stock | | | | |
| cultures. | | 711 | | |
| Chrom Acinetobacter with MDR | MBN0210 | 10 x 90 | 60 | 2-8℃ |
| A Chromogenic media for the isolation of | | mm plates | | |
| Acinetobacter spp. and MDR Acinetobacter | | | | |
| Chromo Candida Agar | MBN0215 | 10 x 90 | 60 | 2-8℃ |
| For detection and differentiation of | | mm plates | | |
| major clinical Candida species. | | | | |
| Chrom Candida Plus | MBN0216 | 10 x 90 | 60 | 2-8℃ |
| For detection and differentiation of | | mm plates | | |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C |
|---------------------------------------------------------------------------------------|----------|----------------------|----------------------|----------------|
| Chrom Campylobacter | MBN0220 | 10 x 90 | 60 | 2-8℃ |
| For detection, differentiation and enumeration of thermotolerant <i>Campylobacter</i> | | mm plates | | |
| Chrom ESBL | MBN0225 | 10 x 90 | 60 | 2-8℃ |
| For the overnight detection of Gram negative bacteria | | mm plates | | |
| producing extended spectrum Beta-Lactamase | | | | |
| Chrom KPC For detection of carbapenem-resistant Enterobacteria (CRE) | MBN0235 | 10 x 90 mm plates | 60 | 2-8℃ |
| Chrom MRSA | | | | |
| For isolation and differentiation of Methicillin | MBN0245 | 10 x 90 | 60 | 2-8℃ |
| Resistant Staphylococcus aureus (MRSA) | | mm plates | | |
| Chrom mSuperCARBA | MBN0247 | 10 x 90 | 60 | 2-8℃ |
| For detection and isolation of Carbapenemase- | | mm plates | | |
| resistant Enterobacteriaceae (CRE) | | | | |
| Chrom O157 For the selective isolation and differentiation of E.coli O157 in | MBN0256 | 10 x 90 | 60 | 2-8℃ |
| food/clinical samples. | | mm plates | | |
| Chrom Salmonella Plus | MBN0265 | 10 x 90 | 60 | 2-8℃ |
| For detection and isolation of Salmonella species | | mm plates | | |
| including lactose positive Salmonella. | | • | | |
| Chrom staph aureus | MBN0270 | 10 x 90 | 60 | 2-8℃ |
| For isolation and direct differentiation of Staphylococcus | | mm plates | | |
| aureus in clinical and industrial samples. | | | | |
| Chrom Strep B | MBN0275 | 10 x 90 | 60 | 2-8℃ |
| For isolation and differentiation of Streptococcus agalactiae | | mm plates | | |
| (GBS) | | | | |
| Chrom UTI | MBN0280 | 10 x 90 | 60 | 2-8℃ |
| For isolation and differentiation of urinary tract pathogens | | mm plates | | |
| Chrom VRE | MBN0285 | 10 x 90 | 60 | 2-8℃ |
| For the detection of Van A / Van B VRE. faecalis & VRE. Faecium | | mm plates | | |
| Chrom Y enterocolitica | MBN0288 | 10 x 90 | 60 | 2-8℃ |
| For detection and differentiation of pathogenic Yersinia | | mm plates | | |
| enterocolitica strains | | | | <u> </u> |
| DCLS Agar (Desoxycholate, Citrate, Lactose, Sucrose) | MBN0290 | 10 x 90 | 60 | 2-8℃ |
| Moderately selective medium for the isolation of Salmonella | | mm plates | | |
| and Shigella from fecal specimens and urine | | 21 | | |
| Desoxycholate Citrate Agar | MBN0291 | 10 x 90 | 60 | 2-8℃ |
| A selective medium for the identification and isolation of | | mm plates | | |
| Salmonella and Shigella from clinical specimens and | | | | |
| environmental samples | | | | |
| DNASE Test Agar | MBN0315 | 10 x 90 | 60 | 2-8℃ |
| A medium for the presumptive identification of pathogenic | | mm plates | | |
| Staphylococci by demonstration of DNase production | | | | |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(|
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|----------------------|----------------------|---------------|
| | • | | | |
| Dermatophyte Test Medium A selective medium for the detection of Dermatophytic fungi | MBN0319 | 10 x 90 mm plates | 90 | 2-8℃ |
| Dichloran Rose Bengal Chloramphenicol (DRBC) Agar Dichloran Rose-Bengal Chloramphenicol Agar (ISO 21527-1) is a selective medium for yeasts and moulds associated with food spoilage. | MBN0325 | 10 x 90 mm plates | 90 | 2-8℃ |
| DG18 agar Dichloran-Glycerol 18% (DG18) (ISO) is a selective medium for yeasts and moulds associated with food spoilage | MBN0330 | 10 x 90 mm plates | 90 | 2-8℃ |
| Diagnostic Sensitivity Test Agar (DST Agar) DST is used as an antibiotic sensitivity-testing medium for antibiotic sensitivity testing of fastidious pathogens such as Neisseria, Streptococcus and Haemophilus species with blood | MBN0331 | 10 x 90 mm plates | 60 | 2-8℃ |
| enrichment from clinical samples. Eosin Methylene Blue Agar (EMB) An isolation medium for the differentiation of the Enterobacteriaceae from clinical, food and environmental | MBN0375 | 10 x 90 mm plates | 90 | 2-8℃ |
| Endo Agar Endo Agar Base is a differential and moderately selective culture medium for the detection and confirmation of coliforms and other enteric microorganisms in waters, milk, dairy and | MBN0378 | 10 x 90 mm plates | 60 | 2-8°C |
| other food products. G. Vaginalis Sel. Medium (V Agar) A medium for the selective isolation of G vaginalis | MBN0422 | 10 x 90 mm plates | 60 | 2-8°C |
| Haemophilus Test Agar Haemophilus Test Agar Base for the susceptibility testing of | MBN0450 | 10 x 90 mm plates | 90 | 2-8°C |
| Haemophilus influenzae Hektoen Enteric Agar A differential and selective medium for the isolation of Salmonella and Shigella from enteric pathological specimens. | MBN0451 | 10 x 90 mm plates | 60 | 2-8℃ |
| Helicobacter Pylori Selective Agar A selective medium for the isolation & cultivation of Helicobacter pylori | MBN0455 | 10 x 90 mm plates | 60 | 2-8℃ |
| Horse Blood agar Horse Blood agar is a highly nutritious general purpose medium for the isolation and cultivation of nonfastidious and | MBN0465 | 10 x 90 mm plates | 60 | 2-8℃ |
| fastidious micro-organisms from clinical specimens. | | | | |
| Hoyle Agar A medium for the isolation and differentiation of Corynebacterium diphtheriae types. | MBN0467 | 10 x 90 mm plates | 60 | 2-8℃ |
| LB agar with Ampicillin Luria Bertani agar by Miller (LB Agar) with Ampicillin is used for the selective cultivation of E. coli strains that contain plasmids conferring ampicillin resistance. | MBN0510 | 10 x 90 mm plates | 60 | 2-8℃ |
| LB agar with Kanamycin | MBN0511 | 10 x 90 | 60 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|----------------------|----------------------|----------------|
| Luria Bertani agar by Miller (LB Agar) with Kanamycin is used for the selective cultivation of E. coli and other organisms that contain plasmids conferring kanamycin resistance. | | mm plates | | |
| Listeria Selective Agar (Oxford) A selective and diagnostic medium for the detection of Listeria monocytogenes from food and clinical laboratories. | MBN0515 | 10 x 90 mm plates | 60 | 2-8℃ |
| Malt Extract Agar with Chloramphenicol Malt extract agar with chloramphenicol is used for the detection, isolation and enumeration of yeasts and moulds. | MBN0530 | 10 x 90 mm plates | 90 | 2-8℃ |
| Martin Lewis Agar | MBN0535 | 10 x 90 | 60 | 2-8°C |
| A selective and enriched medium for the isolation and | | mm plates | | |
| cultivation of Neisseria species from mixed flora | | | | |
| MacConkey Agar w/o Crystal Violet | MBN0540 | 10 x 90 | 90 | 2-8°C |
| A differential medium for the isolation of coliforms and intestinal pathogens in water, dairy products and biological | | mm plates | | |
| specimens. | | | | |
| MacConkey Agar with Crystal Violet A selective medium giving excellent differentiation between coliforms and non-lactose fermenters with inhibition of Gram- | MBN0541 | 10 x 90 mm plates | 90 | 2-8℃ |
| positive micrococci | MENOFAO | 10.00 | 00 | 0.000 |
| MacConkey agar without salt A differential medium on which swarming of Proteus species is suppressed and recommended for urine examination. | MBN0542 | 10 x 90 mm plates | 90 | 2-8℃ |
| MacConkey Agar with Sorbitol | MBN0543 | 10 x 90 | 60 | 2-8℃ |
| A selective and differential medium for the presumptive | | mm plates | | |
| identification of E. Coli O157:h7 | | | | |
| Mannitol Salt Agar A selective medium for the isolation of pathogenic Staphylococci | MBN0547 | 10 x 90 mm plates | 90 | 2-8℃ |
| m-FC Agar m FC Agar is for the detection and enumeration of fecal coliforms by membrane filtration in a laboratory | MBN0550 | 10 x 90 mm plates | 60 | 2-8°C |
| m-Endo Agar m-Endo Agar is used for the enumeration of coliforms in water by the membrane filtration method in a laboratory setting | MBN0556 | 10 x 90 mm plates | 60 | 2-8℃ |
| by the membrane filtration method in a laboratory setting. MILK Plate Count Agar A medium for the bacterial plate counts of microorganisms | MBN0560 | 10 x 90 mm plates | 60 | 2-8°C |
| from milk and dairy derivates | | | | |
| MRS Agar A medium cultivation and enumeration of Lactobacillus species | MBN0565 | 10 x 90 mm plates | 60 | 2-8℃ |
| Mueller Hinton Agar A standardized medium for susceptibility testing by Kirby Bauer | MBN0570 | 10 x 90 mm plates | 90 | 2-8℃ |
| Method Mueller Hinton Blood Agar | MBN0571 | 10 x 90 | 60 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp |
|------------------------------------------------------------------------------|----------|-----------|----------------------|--------------|
| medium for antimicrobial susceptibility testing of | | mm plates | | • |
| Streptococcus pneumoniae | | mm piaces | | |
| Mueller Hinton Chocolate Agar | | | | |
| n apriched modium for the ignicition and cultivation of | MBN0572 | 10 x 90 | 60 | 2-8℃ |
| astidious organisms, particularly Haemophilus species and | MBN0072 | mm plates | 00 | 200 |
| sed for the susceptibility of Haemophilus species | MENIOSOO | <u> </u> | | 0.000 |
| Mycological Agar (deep filled) | MBN0580 | 10 x 90 | 90 | 2-8℃ |
| Mycological agar is a selective medium for the isolation | | mm plates | | |
| f pathogenic fungi, dermatophytes, from clinical specimens. | MENIO | 40.00 | | 0.000 |
| leomycin Blood Agar | MBN0610 | 10 x 90 | 60 | 2-8℃ |
| leomycin blood agar is used for the isolation Clostridia and | | mm plates | | |
| ther anaerobes | | | | |
| lutrient Agar | MBN0615 | 10 x 90 | 90 | 2-8℃ |
| general purpose growth medium for the cultivation of wide | | mm plates | | |
| ariety of bacterial organisms | | 40.00 | | |
| lon nutrient agar | MBN0617 | 10 x 90 | 180 | 2-8℃ |
| medium for the detection of free-living amebae, | | mm plates | | |
| ncluding Acanthamoeba and Naegleria species | | 40.00 | | |
| Oxytetra Glucose Yeast Agar (OGYE Agar) | MBN0625 | 10 x 90 | 90 | 2-8℃ |
| for the selective enumeration of moulds and yeasts. | | mm plates | | |
| Palcam Agar | MBN0630 | 10 x 90 | 60 | 2-8℃ |
| selective and differential diagnostic medium for the detection | | mm plates | | |
| f Listeria monocytogenes | | | | |
| Plate Count Agar | MBN0650 | 10 x 90 | 90 | 2-8℃ |
| medium for the bacterial count in food, water, milk & dairy | | mm plates | | |
| roducts | | | | |
| Potato Dextrose Agar | MBN0655 | 10 x 90 | 90 | 2-8℃ |
| medium for the identification, cultivation and enumeration of | | mm plates | | |
| east and molds in food | | | | |
| Pseudomonas Agar | MBN0660 | 10 x 90 | 60 | 2-8℃ |
| his is a selective medium for the isolation of | | mm plates | | |
| Seudomonas spp. primarily in food, water and | | | | |
| environmental samples. | | | | |
| Seudomonas Cepacia Agar | MBN0663 | 10 x 90 | 60 | 2-8℃ |
| medium for the selective isolation of Bur <mark>kholderia cepacia</mark> | | mm plates | | |
| rom the respiratory secretions of patients with <mark>cystic fibrosis</mark> | | | | |
| nd for routine testing of non-sterile inorganic salt solutions | | | | |
| ontaining preservative. | | | | |
| 2A Agar | | | | |
| 2A agar is used for the enumeration and cultivation of bacteria | MBN0695 | 10 x 90 | 90 | 2-8°C |
| om drinking water | | mm plates | | |
| Rose Bengal Chloramphenicol Agar | MBN0699 | 10 x 90 | 60 | 2-8℃ |
| | | | | |
| BCA the selective enumeration of moulds/yeasts from foods | | mm plates | | |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C |
|------------------------------------------------------------------------------------------------------------------|-------------|-----------|----------------------|----------------|
| Sabouraud Dextrose Agar with Chloramphenicol | MBN0721 | 10 x 90 | 90 | 2-8℃ |
| A medium for the selective isolation of yeast and mould from | MDNO721 | mm plates | | |
| · | | | | |
| food, environmental and clinical samples | MBN0722 | 10 x 90 | 90 | 2-8℃ |
| Sabouraud Dextrose with Chloramphenicol & Gentamicin | WBNOTZZ | mm plates | 00 | 200 |
| A medium for the selective isolation of pathogenic fungi Sabouraud dextrose with Chloramphenicol & Cycloheximide | MBN0723 | 10 x 90 | 90 | 2-8℃ |
| A medium for the selective isolation of pathogenic fungi | IVIDINU123 | mm plates | 90 | 2-0 C |
| Salmonella Shigella Agar | MBN0725 | 10 x 90 | 60 | 2-8℃ |
| A selective medium for the isolation of enteric pathogens | | mm plates | | |
| Schaedler K-V Blood Agar | MBN0730 | 10 x 90 | 60 | 2-8°C |
| A medium for the selective isolation of gram negative anaerobes due to the inclusion of Kanamycin and Vancomycin | | mm plates | | |
| Sheep Blood Agar | MBN0750 | 10 x 90 | 60 | 2-8℃ |
| Ageneral purpose medium for the isolation and cultivation of | | mm plates | | |
| nonfastidious and fastidious microorganisms from clinical and | | | | |
| nonclinical materials Slanetz and Bartley Medium | MBN0753 | 10 x 90 | 60 | 2-8℃ |
| A medium for the detection of enterococci. | IVIDINU133 | mm plates | 00 | 2-0 C |
| Thiosulfate Citrate Bile Salts Sucrose Agar (TCBS) | MBN0780 | 10 x 90 | 60 | 2-8℃ |
| A selective medium for the isolation of Vibrio species | WIDINOTOO | mm plates | 00 | |
| - | MDNIOZOO | • | 00 | 2-8℃ |
| Tryptic Soy Agar (Soyabean- Casein digest agar) | MBN0783 | 10 x 90 | 90 | |
| A general-purpose, non-selective medium capable of | | mm plates | | |
| supporting the growth of most micro-organisms | | | | 2-8℃ |
| Tryptic Soy Agar with Cylcoheximide A medium for the isolation and cultivation of microorganisms | MBN0785 | 10 x 90 | 60 | 2-6 C |
| while inhibiting Saprophytic fungal organisms. | | mm plates | | |
| Perfrigens Agar base (TSC) | MBN0786 | 10 x 90 | 60 | 2-8°C |
| A medium for the detection and enumeration of Clostridium | | mm plates | | |
| Perfringens from food samples | MENIOGO | | 00 | 2-8℃ |
| Violet Red Bile Glucose Agar | MBN0820 | 10 x 90 | 90 | 200 |
| A selective medium for the enterobacteriaceae isolation from | | mm plates | | |
| food, animal feed and environmental samples. | | | - / | 0.0% |
| Violet Red Bile Lactose Agar | MBN0821 | 10 x 90 | 90 | 2-8℃ |
| A lactose-containing selective medium for the detection and | | mm plates | | |
| enumeration of coliform in water, food and dairy products. | | | | |
| VRE Agar | MBN0825 | 10 x 90 | 60 | 2-8℃ |
| A Selective medias for the isolation of Vancomycin Resistant | | mm plates | | |
| Enterococci (VRE) | | | | |
| Wort Agar | MBN0850 | 10 x 90 | 90 | 2-8℃ |
| A medium for the cultivation and enumeration of yeast. | WENTOOO | mm plates | | |
| | MBN0890 | 10 x 90 | 60 | 2-8℃ |
| Xylose Lysine Desoxycholate Agar (XLD) | MDMOOSO | | 00 | |
| A selective medium for the isolation of Salmonellae and | | mm plates | | |
| Shigellae from clinical specimens and foods. | | | | |
| Yeast Extract Glucose Chloramphenicol Agar | MBN0910 | 10 x 90 | 90 | 2-8°C |
| This is a selective medium for the isolation and enumeration of | | mm plates | | |
| yeasts and moulds in dairy products. | | | | |
| Yersinia Selective Agar (CIN Agar) | MBN0915 | 10 x 90 | 60 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C) |
|---------------------------------------------------------|----------|---------------------------|----------------------|-----------------|
| Columbia CNA Blood agar / MacConkey with crystal violet | MBN0950 | 10 x 90 mm Biplates | 60 | 2-8℃ |
| Sheep blood agar / Chocolate agar | MBN0955 | 10 x 90 mm Biplates | 60 | 2-8℃ |
| Sheep blood agar / CLED Agar | MBN0960 | 10 x 90 mm Biplates | 60 | 2-8℃ |
| Sheep blood agar / MacConkey with crystal violet | MBN0970 | 10 x 90 mm Biplates | 60 | 2-8℃ |
| Sheep blood agar / MacConkey without crystal violet | MBN0980 | 10 x 90 mm Biplates | 60 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C) |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|----------------------|----------------------|-----------------|
| Haemophilus Test Agar Haemophilus Test agar for the susceptibility testing of Haemophilus influenzae | MBN1001 | 5 x 150 mm plates | 60 | 2-8℃ |
| Muller Hinton Agar A standardized medium for susceptibility testing by Kirby Bauer Method | MBN1010 | 5 x 150 mm plates | 90 | 2-8°C |
| Mueller Hinton Blood Agar A medium for antimicrobial susceptibility testing of Streptococcus pneumoniae | MBN1015 | 5 x 150 mm plates | 60 | 2-8℃ |
| Sheep Blood Agar A general purpose medium for the isolation and cultivation of nonfastidious and fastidious microorganisms from clinical and non clinical materials | MBN1050 | 5 x 150 mm plates | 60 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp |
|------------------------------------------------------------------------------------------------------------------------------------------------|----------|---------------------|----------------------|--------------|
| Alkaline Peptone Water 5 ml | MBN2101 | 5 ml x 10 | 150 | 2-8℃ |
| A liquid enrichment medium for Vibrio species isolation from | | Tubes | | |
| food, water and clinical samples | | | | |
| Alkaline Peptone Water 10 ml | MBN2102 | 10 ml x 10 | 150 | 2-8℃ |
| A liquid enrichment medium for Vibrio species isolation from | | Tubes | | |
| food, water and clinical samples | | | | |
| Azide Dextrose Broth - Single strength 10 ml Azide Dextrose Broth is used for detection of enterococci in water | MBN2107 | 10 ml x 10 Tubes | 120 | 2-8℃ |
| Azide Dextrose Broth - Double strength 10 ml | MBN2108 | 10 ml x 10 | 120 | 2-8℃ |
| Azide Dextrose Broth is used for detection of enterococci in water | | Tubes | | |
| Brain Heart Infusion (BHI) Broth 5 ML | MBN2125 | 5 ml x 10 | 120 | 2-8℃ |
| Liquid medium for the growth of pathogenic cocci and other | | Tubes | | |
| microorganisms including aerobic and anaerobic bacteria from | | | | |
| a variety of clinical and nonclinical materials | | | | |
| Brain Heart Infusion Agar Slants | MBN2126 | 10 Tubes | 120 | 2-8°C |
| A solid medium which contains the highly nutritious infusions | | | | |
| recommended for the cultivation of fastidious organisms. | | | | |
| Bile aesculin agar slants | MBN2130 | 10 Tubes | 150 | 2-8℃ |
| A differential medium for the isolation and presumptive | | | | |
| dentification of group D streptococci (enterococci) | | | | |
| Brilliant Green Bile Broth (2%) 10 ml | MBN2135 | 10 ml x 10 | 150 | 2-8°C |
| A broth for the detection of coliform bacteria in water, food, | | Tubes | | |
| and dairy products in a laboratory setting | | | | |
| Buffered Peptone Water 10 ml | MBN2140 | 10 ml x 10 | 150 | 2-8℃ |
| Buffered Peptone Water (ISO) is a non-selective pre-enrichment medium for the isolation of Salmonella species from food and associated samples | | Tubes | | |
| Campylobacter enrichment broth (Bolton) 10 ml | MBN2160 | 10 ml x 10 | 90 | 2-8℃ |
| This is a selective enrichment broth for the isolation of | | Tubes | | |
| Campylobacter spp. in clinical, food and environmental | | | | |
| aboratories. | | | | |
| Cooked Meat Broth 10 ml | MBN2161 | 10 ml x 10 | 150 | 2-8℃ |
| A liquid medium for the isolation and cultivation of anaerobic | | Tubes | | |
| and aerobic microorganisms | | | | |
| D/E Neutralizing Broth 9ml | MBN2180 | 9 ml x 10 | 90 | 2-8℃ |
| D/E Neutralizing Broth is used to neutralize and determine the | | Tubes | | |
| pactericidal activity of antiseptics and disinfectants. | | | | |
| E.C broth 10ml | MBN2210 | 10 ml x 10 | 90 | 2-8℃ |
| A selective broth for the growth of coliforms and E. coli from | | Tubes | | |
| food and environmental samples. | | | | |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------|---------------------|----------------------|----------------|
| Fraser Broth 10 ml A secondary selective diagnostic enrichment medium for the isolation of Listeria spp. from food and environmental specimens. | MBN2225 | 10 ml x 10 Tubes | 90 | 2-8℃ |
| Gram negative broth 5 ml For the selective enrichment of Gram-negative microorganisms, especially Shigella spp and Salmonella spp from all types of research materials. | MBN2240 | 5 ml x 10 Tubes | 120 | 2-8℃ |
| Kligler Iron Agar Slant A medium for the identification of Enterobacteriaceae, based on double sugar fermentation and hydrogen sulphide production. | MBN2260 | 10 Tubes | 120 | 2-8℃ |
| Lauryl Sulfate Broth (Lauryl tryptose broth) 10 ml A medium for the detection of coliform organisms in water and waste water, according to the formula of the American Public Health Association | MBN2280 | 10 ml x 10 Tubes | 120 | 2-8℃ |
| Lactose broth 10 ml Lactose broth is used for the performance and confirmation of the Presumptive Test & pre-enrichment medium for coliforms in water and dairy samples. | MBN2281 | 10 ml x 10 Tubes | 120 | 2-8°C |
| Lactose Gelatin Medium 10 ml Lactose Gelatin Medium is used to determine the identity of presumptive Clostridium perfringens. It contains lactose to test lactose fermentation and gelatin to test liquefaction. | MBN2282 | 10 Tubes | 90 | 2-8°C |
| Lim broth 10 ml Lim broth is used for the selective enrichment of group B streptococci | MBN2290 | 10 tubes | 90 | 2-8℃ |
| Loeffler Medium Slants A medium for the cultivation of Corynebacterium diphtheriae from clinical specimens | MBN2295 | 10 Tubes | 120 | 2-8℃ |
| Luria Broth (Miller's LB Broth) 5 ml For molecular genetics studies in E.coli | MBN2305 | 5 ml x 10 Tubes | 150 | 2-8°C |
| Lysine Iron Agar Slants Lysine Iron Agar is used for the differentiation of members of Enterobacteriaceae on the basis of lysine decarboxylase and hydrogen sulfide production. | MBN2310 | 10 Tubes | 150 | 2-8°C |
| Malt extract broth 5 ml Malt extract broth is used for the cultivation of yeasts and moulds and is commonly used as part of sterility testing | MBN2330 | 5 ml x 10 Tubes | 180 | 2-8℃ |
| Maximum Recovery Medium Broth 9 ml A protective and isotonic diluent for maximal recovery of microorganisms | MBN2334 | 9 ml x 10 Tubes | 180 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(0 |
|---------------------------------------------------------------------------------------------------------------------------------------------|-----------|------------|----------------------|----------------|
| Mannitol Selenite Broth 8 ml | MBN2339 | 8 ml x 10 | 150 | 2-8℃ |
| A modification of Selenite F Broth especially recommended for | | Tubes | | |
| the enrichment of salmonellae | | | | |
| MacConkey Broth purple (single strength) 10 ml | MBN2345 | 10 ml x 10 | 180 | 2-8℃ |
| A selective broth for coliforms isolations | | Tubes | | |
| MacConkey Broth purple (double strength) 10 ml | MBN2346 | 10 ml x 10 | 180 | 2-8℃ |
| A selective broth for coliforms isolations | | Tubes | | |
| Motility Test Medium 5 ml | MBN2360 | 5 ml x 10 | 180 | 2-8℃ |
| A medium used for the bacterial motility determination | | Tubes | | |
| MR-VP Test Medium 5 ml | MBN2366 | 5 ml 10 | 180 | 2-8°C |
| A medium for the differentiation of the Escherichia- Enterobacter group (Methyl Red and Voges-Proskauer reactions) from clinical samples | | Tubes | | |
| Mueller Hinton Broth 9 ml | MBN2270 | 9 ml x 10 | 180 | 2-8℃ |
| An antimicrobial susceptibility testing medium which may be | WIDINZZIO | Tubes | 100 | 2-0 0 |
| used in internationally recognized standard procedure | | Tubes | | |
| Mueller-Kanffmann Tetrathonate Novobiocin Broth | MBN2375 | 10 ml x 10 | 60 | 2-8℃ |
| (MkTTn) 10 ml | WIDINZOTO | Tubes | 00 | 2-0 0 |
| A selective enrichment medium for the isolation of Salmonella | | Tubes | | |
| when used with Novobiocin Selective Supplement | | | | |
| Mycological Agar Slant | MBN2380 | 10 Tubes | 180 | 2-8℃ |
| A selective medium for the isolation of pathogenic fungi from | MDMZ300 | 10 Tubes | 100 | 2-0 C |
| materials having a large flora of fungi and bacteria. | | | | |
| 1/500 Nutrient broth 9 ML | MBN2400 | 9 ml x 10 | 180 | 2-8℃ |
| | WDINZ400 | Tubes | 100 | 2-0 C |
| Used for various microbiological applications | MDNIO405 | | 400 | 0.000 |
| Nutrient Agar Slants | MBN2405 | 10 Tubes | 180 | 2-8℃ |
| Nutrient agar slant are primarily used for the cultivation and | | | | |
| maintenance of pure culture | MDNO440 | 5 1 10 | 400 | 0.000 |
| Nutrient Broth 5 ml | MBN2410 | 5 ml x 10 | 180 | 2-8℃ |
| For the cultivation of non-fastidious microorganisms in water, | | Tubes | | |
| feces and other materials | | | | |
| Nutrient Broth 9 ml | MBN2411 | 9 ml x 10 | 180 | 2-8℃ |
| For the cultivation of non-fastidious microorganisms in water, | | Tubes | | |
| feces and other materials | AT. | 1 | | |
| Peptone water 5ml | MBN2450 | 5 ml x 10 | 180 | 2-8℃ |
| Peptone water is a basal medium to which carbohydrates and | | Tubes | | |
| indicator may be added for fermentation studies. | | | | |
| Listeria Enrichment Broth UVM II - Secondary 10 ml | MBN2461 | 10 ml x 10 | | |
| A secondary selective enrichment supplement for the isolation | | Tubes | 120 | 2-8℃ |
| of listeria when used with Listeria Enrichment Broth Base | | | | |
| Phosphate buffered saline (PBS) 10 ML | MBN2490 | 10 ml x 10 | 180 | 2-8℃ |
| PBS is a buffer solutions commonly used in biological research | | Tubes | | |
| Rappaport Vassiliadis Soy Broth (RVS broth) 10 ml | MBN2510 | 10 ml x 10 | 180 | 2-8°C |
| An enrichment medium for the isolation of Salmonella species | | Tubes | | |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(|
|---------------------------------------------------------------------------------------------------------------------------------------------|----------|------------|----------------------|---------------|
| Mannitol Selenite Broth 8 ml | MBN2339 | 8 ml x 10 | 150 | 2-8℃ |
| A modification of Selenite F Broth especially recommended for | | Tubes | | |
| the enrichment of salmonellae | | | | |
| MacConkey Broth purple (single strength) 10 ml | MBN2345 | 10 ml x 10 | 180 | 2-8℃ |
| A selective broth for coliforms isolations | | Tubes | | |
| MacConkey Broth purple (double strength) 10 ml | MBN2346 | 10 ml x 10 | 180 | 2-8℃ |
| A selective broth for coliforms isolations | | Tubes | | |
| Motility Test Medium 5 ml | MBN2360 | 5 ml x 10 | 180 | 2-8°C |
| A medium used for the bacterial motility determination | | Tubes | | |
| MR-VP Test Medium 5 ml | MBN2366 | 5 ml 10 | 180 | 2-8℃ |
| A medium for the differentiation of the Escherichia- Enterobacter group (Methyl Red and Voges-Proskauer reactions) from clinical samples | | Tubes | | |
| Mueller Hinton Broth 9 ml | MBN2270 | 9 ml x 10 | 180 | 2-8℃ |
| An antimicrobial susceptibility testing medium which may be | | Tubes | 100 | _ 0 0 |
| used in internationally recognized standard procedure | | 14500 | | |
| Mueller-Kanffmann Tetrathonate Novobiocin Broth | MBN2375 | 10 ml x 10 | 60 | 2-8℃ |
| (MkTTn) 10 ml | | Tubes | | |
| A selective enrichment medium for the isolation of Salmonella | | | | |
| when used with Novobiocin Selective Supplement | | | | |
| Mycological Agar Slant | MBN2380 | 10 Tubes | 180 | 2-8℃ |
| A selective medium for the isolation of pathogenic fungi from | | | | |
| materials having a large flora of fungi and bacteria. | | | | |
| 1/500 Nutrient broth 9 ML | MBN2400 | 9 ml x 10 | 180 | 2-8℃ |
| Used for various microbiological applications | | Tubes | | |
| Nutrient Agar Slants | MBN2405 | 10 Tubes | 180 | 2-8℃ |
| Nutrient agar slant are primarily used for the cultivation and | | | | |
| maintenance of pure culture | | | | |
| Nutrient Broth 5 ml | MBN2410 | 5 ml x 10 | 180 | 2-8℃ |
| For the cultivation of non-fastidious microorganisms in water, | | Tubes | | |
| feces and other materials | | | | |
| Nutrient Broth 9 ml | MBN2411 | 9 ml x 10 | 180 | 2-8℃ |
| For the cultivation of non-fastidious microorganisms in water, | | Tubes | | |
| feces and other materials | TA | 1 | | |
| Peptone water 5ml | MBN2450 | 5 ml x 10 | 180 | 2-8℃ |
| Peptone water is a basal medium to which carbohydrates and | | Tubes | | |
| indicator may be added for fermentation studies. | | 411 | | |
| Listeria Enrichment Broth UVM II - Secondary 10 ml | MBN2461 | 10 ml x 10 | | |
| A secondary selective enrichment supplement for the isolation | | Tubes | 120 | 2-8℃ |
| of listeria when used with Listeria Enrichment Broth Base | | | | |
| Phosphate buffered saline (PBS) 10 ML | MBN2490 | 10 ml x 10 | 180 | 2-8℃ |
| PBS is a buffer solutions commonly used in biological research | | Tubes | | |
| Rappaport Vassiliadis Soy Broth (RVS broth) 10 ml | MBN2510 | 10 ml x 10 | 180 | 2-8℃ |
| An enrichment medium for the isolation of Salmonella species | | Tubes | | |

| Rappaport Vassiliadis Broth (RV) 10 ML | • | | | |
|---------------------------------------------------------------------|---------|------------|-----|-------|
| | | | | |
| | MBN2512 | 10 ml x 10 | 180 | 2-8℃ |
| A selective enrichment broth for the isolation of salmonellae | | Tubes | | |
| Ringer solutions 1/4 strength 9 ml | MBN2521 | 9 ml x 10 | 270 | 2-8℃ |
| An osmotically controlled solution for the preparation of | | Tubes | | |
| suspensions of food samples and for use as a diluent in dilution | | | | |
| techniques for bacterial enumeration | | | | |
| Saline normal 5 ml | MBN2540 | 5 ml x 10 | 270 | 2-8°C |
| Commonly used in Laboratory as diluting reagents | | Tubes | | |
| Salt broth 6.5% 5 ml | MBN2541 | 5 ml x 10 | 270 | 2-8°C |
| Salt Broth, Modified is used for cultivation and differentiation of | | Tubes | | |
| the enterococcal group D streptococci from nonenterococcal | | | | |
| group D streptococci based on salt tolerance | | | | |
| Sabouraud Dextrose Agar Slants | MBN2550 | 10 Tubes | 180 | 2-8℃ |
| A medium for the cultivation of fungi | | | | |
| Sabouraud Dextrose Broth 5 ml | MBN2555 | 5 ml x 10 | 180 | 2-8℃ |
| Sabouraud Dextrose Broth is used for cultivation of yeasts, | | Tubes | | |
| moulds and aciduric microorganisms | | | | |
| Salmonella Shigella Broth 5 ml | MBN2557 | 5 ml x 10 | 150 | 2-8℃ |
| For the cultivation of Salmonella and Shigella | | Tubes | | |
| SCDLP Broth 10 ml | MBN2560 | 10 ml x 10 | 60 | 2-8°C |
| Measurement of antibacterial activity on plastics surfaces as | | Tubes | | |
| per ISO 22196 | | | | |
| Selenite Cystine Broth 8 ml | MBN2570 | 8 ml x 10 | 150 | 2-8℃ |
| A selective enrichment of Salmonella species and some strains | | Tubes | | |
| of Shigella in feces, urine | | | | |
| Selenite F Broth 8 ml | MBN2571 | 8 ml x 10 | 150 | 2-8℃ |
| A Selective enrichment of Salmonella species and some strains | | Tubes | | |
| of Shigella in feces, urine | | | 1 | |
| Simmons Citrate Slant | MBN2577 | 10 Tubes | 180 | 2-8℃ |
| A medium for the determination of citrate utilization of | | | | |
| Enterobacteriaceae | | | | |
| SIM Medium 5 ml | MBN2578 | 5 ml x 10 | 150 | 2-8℃ |
| A medium for the identification and differentiation of | | Tubes | | |
| Enterobacteriaceae | | mi/ | | |
| Tetrathionate broth 10 ml | MBN2600 | 10 ml x 10 | 90 | 2-8°C |
| Tetrathionate Broth is recommended for the selective | | Tubes | | |
| enrichment method of isolating Salmonella typhi and other | | | | |
| salmonellae from faeces, sewage, etc. | | | | |
| Thioglycollate broth 10 ml | MBN2610 | 10 ml x 10 | 150 | 2-8℃ |
| A medium for the cultivation of both aerobic and anaerobic | | Tubes | | |
| organisms in the performance of sterility tests | | | | |
| Triple Sugar Iron Agar Slant | | | | |
| A medium for the differentiation of Enterobacteriaceae based | MBN2615 | 10 Tubes | 180 | 2-8°C |

| on hydrogen sulphide production and fermentation of lactose, sucrose and dextrose | | | | |
|----------------------------------------------------------------------------------------------------------------------------------------|---------|---------------------|-----|------|
| Tryptone Soy Agar Slants A general purpose medium for multiple application including culture of fastidious bacteria | MBN2618 | 10 Tubes | 180 | 2-8℃ |
| Trypticase Soy Broth 10 ml A highly nutritious general purpose medium for the growth of bacteria and fungi | MBN2619 | 10 ml x 10 Tubes | 180 | 2-8℃ |
| Trypticase Soy Broth 5 ml A highly nutritious general purpose medium for the growth of bacteria and fungi | MBN2620 | 5 ml x 10 Tubes | 180 | 2-8℃ |
| Tryptone Water 5 ml A liquid medium for the production of indole by micro- organisms. | MBN2621 | 5 ml x 10 Tubes | 180 | 2-8℃ |
| Trypticasein Soy Broth with Novobiocine (mTSB) 10 ml For the selective enrichment of Escherichia coli O157 from food or faecal samples | MBN2625 | 10 ml x 10 Tubes | 120 | 2-8℃ |
| Urea Agar Slants A medium for the confirmation of Enterobacteriaceae on the basis of urease production. | MBN2650 | 10 Tubes | 180 | 2-8℃ |
| Water sterile 5 ML Sterile water is free of microorganisms such as bacteria, fungi spores and is used in laboratory. | MBN2680 | 5 ml x 10 Tubes | 365 | 2-8℃ |

| Product Name | REF Code | Size | Shelf Life (days) | Storage Temp(C |
|---------------------------------------------------------------------------------------------------|----------|---------------|----------------------|----------------|
| Alkalina Pantana Water | MDNI2400 | 400 1 | 450 | 2.0% |
| Alkaline Peptone Water | MBN3100 | 100 ml bot | 150 | 2-8℃ |
| A liquid enrichment medium for vibrio species isolation from food, water and clinical samples | | DOL | | |
| Plate Count Agar | MBN3140 | 100 ml | 90 | 2-8℃ |
| A medium for the bacterial count in food, water, milk and other | W.D. (6) | bot | | 200 |
| dairy products | | | | |
| Violet Red Bile Lactose Agar (VRBLA) | MBN3290 | 100 ml | 90 | 2-8℃ |
| A lactose-containing selective medium for the detection and | | bot | | |
| enumeration of coliform in water, food and dairy products | | | | |
| Alkaline Peptone Water | MBN4100 | 250 ml | 150 | 2-8℃ |
| A liquid enrichment medium for Vibrio species isolation | | bot | | |
| EE Broth (Buffered brilliant bile broth) | MBN4110 | 250 ml | 120 | 2-8℃ |
| An enrichment medium for Enterobacteriaceae in the | | bot | | |
| bacteriological examination of foods. | | | | |
| Half Fraser Broth 225 ML | MBN4130 | 250 ml | 150 | 2-8℃ |
| A selective supplement for the primary enrichment of Listeria | | bot | | |
| spp. from food and environmental samples when used with | | | | |
| Fraser Broth | | | | |
| Listeria Enrichment Broth UVM I - Primary 225 ML | MBN4150 | 250 ml | 120 | 2-8℃ |
| A primary selective enrichment supplement for the isolation of | | bot | | |
| listeria when used with Listeria Enrichment Broth Base | | | | |
| Orange serum agar | MBN4180 | 250 ml | 90 | 2-8℃ |
| A medium for the isolation and enumeration of spoilage | | bot | | |
| organisms of citrus products. | | | | |
| One Broth Salmonella | MBN4181 | 250 ml | 90 | 2-8℃ |
| A selective medium for the recovery of Salmonella spp.in a | | bot | | |
| single enrichment step | MENIAGAG | 050 | 00 | 0.000 |
| Plate Count Agar | MBN4210 | 250 ml | 90 | 2-8℃ |
| A medium for the bacterial count in food, water, milk and other | | bot | | |
| dairy products | MPNIAGO | 250 ml | 90 | 2-8℃ |
| Violet Red Bile Lactose Agar (VRBLA) A lactose-containing selective medium for the detection and | MBN4260 | bot | 90 | 2-6 C |
| enumeration of coliform in water, food and dairy products | | DOL | | |
| | MPNE100 | 500 ml | 150 | 2 000 |
| Aliquid enrichment medium for Vibrio species isolation | MBN5100 | 500 ml | 150 | 2-8℃ |
| A liquid enrichment medium for Vibrio species isolation | | bot | | |

